

Mth100 Past papers for mid term

$$A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 1 & 2 \\ 3 & 1 & 3 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} 2 & 2 & 3 \\ 2 & 1 & 1 \end{bmatrix}$$

Check whether the matrix A and B can be multiplied? Justify your answer.


Q

Solve following system of equations:

$$2x_1 + x_2 + 3x_3 = 11$$

$$4x_1 - x_2 + 2x_3 = 5$$

$$3x_2 + 2x_3 = 13$$



Created by:

Hafiza Mubeen, Muhammad Nouman, Tasha Khan, M Qamar Nazir Page 13

Mth100 Past papers for mid term

Mth100 Past papers for mid term

There are five juniors and three seniors in the Service Club. The club is to send four representatives to the State Conference. If the members of the club decide to send two juniors and two seniors, how many different groupings are possible?

Q

$$A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 1 & 2 \\ 0 & 0 & 1 \end{bmatrix}, I = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

If $A^2 - 5I$ then find A^{-1}

Q

Find the 10th term of the geometric sequence with $a = 3000$ and a common ratio $r = \frac{1}{2}$

Q

Let $f(x) = 3x + 1$ and $g(x) = 2x + 3$. Find the product of f & g i.e. $(f \cdot g)(x)$

Q

If $a_1 = 2, a_2 = 3, a_3 = -2,$ and $a_4 = 1$, then compute the summation: $\sum_{i=1}^4 2a_i$?

Q

The local Family Restaurant has a daily breakfast special in which the customer may choose one item from each of the following groups:

Breakfast Sandwich	Accompaniments	Juice
egg and chicken	breakfast potatoes	orange
egg and beef	apple slices	cranberry
egg and cheese	fresh fruit cup	tomato
	pastry	apple
		grape

How many different breakfast specials are possible?

Q

Created by:

Hafiza Mubeen, Muhammad Nouman, Tasha Khan, M Qamar Nazir Page 12

Mth100 Past papers for mid term

$$A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 1 & 2 \end{bmatrix}$$

Mth100 Past papers for mid term

b) $r_n = \frac{r_1^n}{(n-r)!}$

c) $r_n = \frac{r_1^n}{(n-r)!r!}$

d) $r_n = r_1^n (n-r)!$

Q

det

a) -10

b) 10

c) -2

d) -2

Q

An Arithmetic Sequence is a sequence in which each term after the first term is found by _____ a constant called *common difference*.

- a) Adding
- b) Subtracting
- c) Dividing
- d) Multiplying

Q

To multiply a matrix by a scalar, we multiply each _____ in the matrix by that scalar.

- a) Row
- b) Column
- c) Entry
- d) Vector

Q

Write the domain and range of $f(x) = 5x - 2$

Q

List all the missing terms in the arithmetic series: 7, 12, 17, ..., 37

Q

Solve the equation $x^2 - 6x - 5 = 0$ by factorization method

Q

Created by:

Hafiza Mubeen, Muhammad Nouman, Tasha Khan, M Qamar Nazir Page 11

Mth100 Past papers for mid term

There are five juniors and three seniors in the Service Clubs. The club is to send four representatives to the State Conference. If the members of the club decide to send two juniors and two seniors, how many different groupings are possible?

Mth100 Past papers for mid term

d) $f(x) \geq 3$

Q

If a scalar k is multiplied with the matrix then which one of the following statement is correct?

- a) k will be multiplied with all elements of the matrix
- b) k will be multiplied with first column in the matrix
- c) k will be multiplied with first row in the matrix
- d) k will be multiplied with the diagonal elements

Q

Let ' a ' be the first term and ' d ' be the common difference of an arithmetic series. Then

which of the following is the sum (S_n) of the first ' n ' terms of an arithmetic series?

- a) $a + (n-1)d$
- b) $2a + (n-1)d$
- c) $\frac{n}{2}[a + (n-1)d]$
- d) $\frac{n}{2}[2a + (n-1)d]$

Q

A Square matrix with ones on the diagonal and zeros elsewhere is called an matrix
rectangular
adjoint
determinant
identity

Q

Which of the following is true for nC_r ?

a) ${}^nC_r = \frac{{}^nP_r}{r!}$

Created by:

Hafiza Mubeen, Muhammad Nouman, Tasha Khan, M Qamar Nazir Page 10

Mth100 Past papers for mid term

c) $I - \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$

d) $I - \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

Q

Let $z_1 = (3 + 5i)$ and $z_2 = (-9 + 6i)$ be two complex numbers, then which of the following is the value of $\frac{z_1}{z_2}$?

- a) $-6-i$
- b) $-6+i$
- c) $6-i$
- d) $6+i$

Q

A sequence, in which each term after the first is found by multiplying the previous term by a constant value, is called a (an) _____.

- a) Arithmetic series
- b) Arithmetic progression
- c) Geometric series
- d) Geometric progression

Q

Which of these relations is function?

- a) One-many relation
- b) Many-one relation
- c) Many-many relation
- d) None of these

Q

The range of $f(x) = (x - 2)^2 + 3$ is.....

- a) $f(x) > 3$
- b) $f(x) < 3$
- c) $f(x) \geq 3$

Created by:

Hafiza Mubeen, Muhammad Nouman, Tasha Khan, M Qamar Nazir Page 9

Mth100 Past papers for mid term

$A = \{2, 4, 6, 8, 10, 12, 14, 16\}$ and $B = \{12, 13, 14, 15, 16\}$
If $A \cap B$ is

- a) $A \cup B = \{2, 4, 6, 8, 10, 12, 13, 14, 15, 16\}$
- b) $A \cup B = \{12, 13, 14, 15, 16\}$
- c) $A \cap B = \{12, 14, 16\}$
- d) $A \cap B = \{2, 4, 6, 8, 10, 12, 14, 16\}$

Q
5.45235452365845125... is a/an

- a) rational number
- b) irrational number
- c) integer
- d) natural number

Q
The inverse of the non-singular matrix A can be found by the formula

- a) $A^{-1} = A \text{adj}(A)$
- b) $A^{-1} = \frac{1}{A} \text{adj}(A)$
- c) $A^{-1} = \frac{1}{|A|} \text{adj}(A)$
- d) $A^{-1} = \frac{1}{|A^{-1}|} \text{adj}(A)$

Q
Which of the following is identity matrix

- a) $I = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$
- b) $I = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$

Created by:

Hafiza Mubeen, Muhammad Nouman, Tasha Khan, M Qamar Nazir Page 8

Mth100 Past papers for mid term

- c) 6, 12, 20, ...
- d) 12, 20, 30, ...

Q

If a sequence of numbers approaches to a finite number, then which of the following is true?

- a) The sequence is infinite
- b) The sequence is divergent
- c) The sequence is convergent
- d) None of them is true.

Q

A function of the type $y = ax^2 + bx + c$ where a , b and c are real coefficients, is called a

- a) Quadratic function
- b) Linear Function
- c) Constant function
- d) Cubic Function

Q

If the slopes of two lines are same then these lines are

- a) Parallel
- b) Perpendicular
- c) Not parallel
- d) None of these

Q

Determine the inverse of the function $f(x) = 4x^2 - 4$

- a) $f^{-1}(x) = \frac{\sqrt{x+4}}{4}$
- b) $f^{-1}(x) = \frac{\sqrt{x-4}}{4}$
- c) $f^{-1}(x) = \frac{\sqrt{x+4}}{2}$
- d) $f^{-1}(x) = \frac{\sqrt{x-4}}{2}$

Q

Created by:

Hafiza Mubeen, Muhammad Nouman, Tasha Khan, M Qamar Nazir Page 7

Mth100 Past papers for mid term

d) identity

Q

If the order does matter in a set of objects, it is a _____.

- a) sequence
- b) combination
- c) permutation
- d) sample

Q

The Minor of an element in a third-order determinant is a _____ determinant obtained by deleting the row and column that contains the element.

- a) third order
- b) second order
- c) fourth order
- d) fifth order

Q

$$A = 2 \begin{bmatrix} -2 & 0 & 1 \\ 4 & -1 & 3 \end{bmatrix}$$

Consider _____ be a scalar matrix then which of the following is the dimension of the matrix A ?

- a) 3×2
- b) 2×1
- c) 2×2
- d) 2×3

Q

Which of these equations has two roots?

- a) Quadratic
- b) Linear
- c) Cubic
- d) None of these

Q

$$C_n = n(n+1)$$

If _____ for all integers $n \geq 2$, then which of the following is the first three terms of a sequence?

- a) 0, 2, 6, ...
- b) 2, 6, 12, ...

Created by:

Hafiza Mubeen, Muhammad Nouman, Tasha Khan, M Qamar Nazir Page 6

Mth100 Past papers for mid term

- b) $\frac{1}{2}$
c) 2
d) 5

Q

If $f(x) = 2x + 1$, $g(x) = x^2 - 1$ then $f \circ g(x) =$

- a) $2x^2$
b) $2x^2 - 1$
c) $2x^2 + 1$
d) $2x - 1$

Q

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

For _____, there will be exactly one real root if

- a) $b^2 - 4ac > 0$
b) $b^2 - 4ac \leq 0$
c) $b^2 - 4ac = 0$
d) $b^2 - 4ac < 0$

Q

If A is a matrix whose determinant is zero then its inverse _____.

- a) will be zero
b) cannot be determined
c) will be negative
d) will be positive

Q

Which of the following is true for the list $2 + 4 + 6 + 8 + 10$?

- a) This is an arithmetic sequence.
b) This is a geometric sequence.
c) This is an arithmetic series.
d) This is a geometric series.

Q

A Square matrix with ones on the diagonal and zeros elsewhere is called an _____ matrix

- a) rectangular
b) adjoint
c) determinant

Created by:

Hafiza Mubeen, Muhammad Nouman, Tasha Khan, M Qamar Nazir Page 5

Mth100 Past papers for mid term

- d) identity

Q

If the order does matter in a set of objects, it is a _____.

- a) sequence
b) combination

Mth100 Past papers for mid term

- h) 10
- c) 14
- d) 8

Q

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$

Let $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$. The Minor of a_{23} is

- a) $a_{11} \ a_{12}$
- b) $\begin{vmatrix} a_{11} & a_{12} \\ a_{31} & a_{32} \end{vmatrix}$
- c) $\begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix}$
- d) $\begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix}$

Q

For an arithmetic sequence $a, a + (n-1)d$, which of the following is the best option for 'd'?

- a) First term
- b) Last term
- c) Constant
- d) Common difference

Q

Which of the following is the common ratio 'r' of an geometric sequence 5, 10, 20, 40, ...?

- a) -8

Created by:

Hafiza Mubeen, Muhammad Nouman, Tasha Khan, M Qamar Nazir Page 4

Mth100 Past papers for mid term

Q4

Let $z_1 = 3 - 5i$ and $z_2 = -4 - 2i$ be two complex numbers, then which of the following is

the value of $z_1 \cdot z_2$?

- a) $-22 + 14i$
- b) $-12 + 10i$
- c) $-6 + 20i$
- d) $14 - 22i$

Q

Let $z_1 = 2 + 3i$ and $z_2 = 5 + 9i$ be two complex numbers, then which of the following is

the value of $z_1 + z_2$?

- a) $7 - 12i$
- b) $7 + 12i$
- c) $10 + 27i$
- d) $18 + 15i$

Q

What is roster method to show the set $A = \{x | x \in \mathbb{Z} \wedge 3 < x \leq 3\}$?

- a) $A = \{-3, -2, -1, 0, 1, 2, 3\}$
- b) $A = \{-2, -1, 0, 1, 2, 3\}$
- c) $A = \{-3, -2, -1, 0, 1, 2\}$
- d) $A = \{-3, -2, -1, 1, 2, 3\}$

Q

5.45235452365845125... is a/an

- a) rational number
- b) irrational number
- c) integer
- d) natural number

Q

$$A = \begin{bmatrix} -1 & 2 \\ -3 & -4 \end{bmatrix}$$

Determinant of A is

- a) 12

Created by:

Hafiza Mubeen, Muhammad Nouman, Tasha Khan, M Qamar Nazir Page 3

Mth100 Past papers for mid term

- b) 10
- c) 11
- d) 9

Mth100 Past papers for mid term

c) $I - \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$

d) $I - \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

Q

Let $z_1 = (3 + 5i)$ and $z_2 = (-9 - 6i)$ be two complex numbers, then which of the following is the value of $\frac{z_1}{z_2}$?

- a) $-6-i$
- b) $-6+i$
- c) $6-i$
- d) $6+i$

Q

A sequence, in which each term after the first is found by multiplying the previous term by a constant value, is called a (an) _____.

- a) Arithmetic series
- b) Arithmetic progression
- c) Geometric series
- d) Geometric progression

Q

Which of these relations is function?

- a) One-many relation
- b) Many-one relation
- c) Many-many relation
- d) None of these

Q

The range of $f(x) = (x - 2)^2 + 3$ is.....

- a) $f(x) > 3$
- b) $f(x) < 3$
- c) $f(x) \geq 3$

Created by:

Hafiza Mubeen, Muhammad Nouman, Tasha Khan, M Qamar Nazir Page 9

Mth100 Past papers for mid term

$A = \{2, 4, 6, 8, 10, 12, 14, 16\}$ and $B = \{12, 13, 14, 15, 16\}$
If $A \cap B$ is

- a) $A \cup B = \{2, 4, 6, 8, 10, 12, 13, 14, 15, 16\}$
- b) $A \cup B = \{12, 13, 14, 15, 16\}$
- c) $A \cap B = \{12, 14, 16\}$
- d) $A \cap B = \{2, 4, 6, 8, 10, 12, 14, 16\}$

Q

5.45235452365845125... is a/an

- a) rational number
- b) irrational number
- c) integer
- d) natural number

Q

The inverse of the non-singular matrix A can be found by the formula

- a) $A^{-1} = A \text{adj}(A)$
- b) $A^{-1} = \frac{1}{A} \text{adj}(A)$
- c) $A^{-1} = \frac{1}{|A|} \text{adj}(A)$
- d) $A^{-1} = \frac{1}{|A^{-1}|} \text{adj}(A)$

Q

Which of the following is identity matrix

- a) $I = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$
- b) $I = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$

Created by:

Hafiza Mubeen, Muhammad Nouman, Tasha Khan, M Qamar Nazir Page 8

Mth100 Past papers for mid term

- c) 6, 12, 20, ...
- d) 12, 20, 30, ...

Q

If a sequence of numbers approaches to a finite number, then which of the following is true?

- a) The sequence is infinite
- b) The sequence is divergent
- c) The sequence is convergent
- d) None of them is true.

Q

A function of the type $y = ax^2 + bx + c$ where a , b and c are real coefficients, is called a

- a) Quadratic function
- b) Linear Function
- c) Constant function
- d) Cubic Function

Q

If the slopes of two lines are same then these lines are

- a) Parallel
- b) Perpendicular
- c) Not parallel
- d) None of these

Q

Determine the inverse of the function $f(x) = 4x^2 - 4$

a) $f^{-1}(x) = \frac{\sqrt{x+4}}{4}$

b) $f^{-1}(x) = \frac{\sqrt{x-4}}{4}$

c) $f^{-1}(x) = \frac{\sqrt{x+4}}{2}$

d) $f^{-1}(x) = \frac{\sqrt{x-4}}{2}$

Q

Created by:

Hafiza Mubeen, Muhammad Nouman, Tasha Khan, M Qamar Nazir Page 7

Mth100 Past papers for mid term

d) identity

Q

If the order does matter in a set of objects, it is a _____.

- a) sequence
- b) combination
- c) permutation
- d) sample

Q

The Minor of an element in a third-order determinant is a _____ determinant obtained by deleting the row and column that contains the element.

- a) third order
- b) second order
- c) fourth order
- d) fifth order

Q

$$A = 2 \begin{bmatrix} -2 & 0 & 1 \\ 4 & -1 & 3 \end{bmatrix}$$

Consider _____ be a scalar matrix then which of the following is the dimension of the matrix A ?

- a) 3×2
- b) 2×1
- c) 2×2
- d) 2×3

Q

Which of these equations has two roots?

- a) Quadratic
- b) Linear
- c) Cubic
- d) None of these

Q

$$C_n = n(n+1)$$

If _____ for all integers $n \geq 2$, then which of the following is the first three terms of a sequence?

- a) 0, 2, 6, ...
- b) 2, 6, 12, ...

Created by:

Hafiza Mubeen, Muhammad Nouman, Tasha Khan, M Qamar Nazir Page 6

Mth100 Past papers for mid term

- b) $\frac{1}{2}$
c) 2
d) 5

Q

If $f(x) = 2x + 1$, $g(x) = x^2 - 1$ then $f \circ g(x) =$

- a) $2x^2$
b) $2x^2 - 1$
c) $2x^2 + 1$
d) $2x - 1$

Q

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

For _____, there will be exactly one real root if

- a) $b^2 - 4ac > 0$
b) $b^2 - 4ac \leq 0$
c) $b^2 - 4ac = 0$
d) $b^2 - 4ac < 0$

Q

If A is a matrix whose determinant is zero then its inverse _____.

- a) will be zero
b) cannot be determined
c) will be negative
d) will be positive

Q

Which of the following is true for the list $2 + 4 + 6 + 8 + 10$?

- a) This is an arithmetic sequence.
b) This is a geometric sequence.
c) This is an arithmetic series.
d) This is a geometric series.

Q

A Square matrix with ones on the diagonal and zeros elsewhere is called an _____ matrix

- a) rectangular
b) adjoint
c) determinant

Created by:

Hafiza Mubeen, Muhammad Nouman, Tasha Khan, M Qamar Nazir Page 5

Mth100 Past papers for mid term

- d) identity

Q

If the order does matter in a set of objects, it is a _____.

- a) sequence
b) combination

Mth100 Past papers for mid term

- h) 10
- c) 14
- d) 8

Q

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$

Let $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$. The Minor of a_{23} is

- a) $a_{11} \ a_{12}$
- b) $\begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix}$
- c) $\begin{vmatrix} a_{11} & a_{12} \\ a_{31} & a_{32} \end{vmatrix}$
- d) $\begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix}$

Q For an arithmetic sequence $a, a + (n-1)d$, which of the following is the best option for 'd'?

- a) First term
- b) Last term
- c) Constant
- d) Common difference

Q Which of the following is the common ratio 'r' of an geometric sequence 5, 10, 20, 40, ...?

- a) -8

Created by:

Hafiza Muheeb, Muhammad Nouman, Tasha Khan, M Qamar Nazir Page 4

Mth100 Past papers for mid term

Q4

Let $z_1 = 3 - 5i$ and $z_2 = -4 - 2i$ be two complex numbers, then which of the following is

the value of $z_1 \cdot z_2$?

- a) $-22 + 14i$
- b) $-12 + 10i$
- c) $-6 + 20i$
- d) $14 - 22i$

Q

Let $z_1 = 2 + 3i$ and $z_2 = 5 + 9i$ be two complex numbers, then which of the following is

the value of $z_1 + z_2$?

- a) $7 - 12i$
- b) $7 + 12i$
- c) $10 + 27i$
- d) $18 + 15i$

Q

What is roster method to show the set $A = \{x | x \in \mathbb{Z} \wedge 3 < x \leq 3\}$?

- a) $A = \{-3, -2, -1, 0, 1, 2, 3\}$
- b) $A = \{-2, -1, 0, 1, 2, 3\}$
- c) $A = \{-3, -2, -1, 0, 1, 2\}$
- d) $A = \{-3, -2, -1, 1, 2, 3\}$

Q

5.45235452365845125... is a/an

- a) rational number
- b) irrational number
- c) integer
- d) natural number

Q

$$A = \begin{bmatrix} -1 & 2 \\ -3 & -4 \end{bmatrix}$$

Determinant of A is

- a) 12

Created by:

Hafiza Mubeen, Muhammad Nouman, Tasha Khan, M Qamar Nazir Page 3

Mth100 Past papers for mid term

- b) 10
- c) 11
- d) 9

Mth100 Past papers for mid term

- b) $\frac{1}{2}$
c) 2
d) 5

Q

If $f(x) = 2x + 1$, $g(x) = x^2 - 1$ then $f \circ g(x) =$

- a) $2x^2$
b) $2x^2 - 1$
c) $2x^2 + 1$
d) $2x - 1$

Q

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

For _____, there will be exactly one real root if

- a) $b^2 - 4ac > 0$
b) $b^2 - 4ac \leq 0$
c) $b^2 - 4ac = 0$
d) $b^2 - 4ac < 0$

Q

If A is a matrix whose determinant is zero then its inverse _____.

- a) will be zero
b) cannot be determined
c) will be negative
d) will be positive

Q

Which of the following is true for the list $2 + 4 + 6 + 8 + 10$?

- a) This is an arithmetic sequence.
b) This is a geometric sequence.
c) This is an arithmetic series.
d) This is a geometric series.

Q

A Square matrix with ones on the diagonal and zeros elsewhere is called an _____ matrix

- a) rectangular
b) adjoint
c) determinant

Created by:

Hafiza Mubeen, Muhammad Nouman, Tasha Khan, M Qamar Nazir Page 5

Mth100 Past papers for mid term

- d) identity

Q

If the order does matter in a set of objects, it is a _____.

- a) sequence
b) combination

Mth100 Past papers for mid term

- h) 10
- c) 14
- d) 8

Q

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$

Let $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$. The Minor of a_{23} is

- a) $a_{11} \ a_{12}$
- b) $\begin{vmatrix} a_{11} & a_{12} \\ a_{31} & a_{32} \end{vmatrix}$
- c) $\begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix}$
- d) $\begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix}$

Q For an arithmetic sequence $a, a+(n-1)d$, which of the following is the best option for 'd'?

- a) First term
- b) Last term
- c) Constant
- d) Common difference

Q Which of the following is the common ratio 'r' of an geometric sequence 5, 10, 20, 40, ...?

- a) -8

Created by:

Hafiza Mubeen, Muhammad Nouman, Tasha Khan, M Qamar Nazir Page 4

Mth100 Past papers for mid term

Q4

Let $z_1 = 3 - 5i$ and $z_2 = -4 - 2i$ be two complex numbers, then which of the following is

the value of $z_1 \cdot z_2$?

- a) $-22 + 14i$
- b) $-12 + 10i$
- c) $-6 + 20i$
- d) $14 - 22i$

Q

Let $z_1 = 2 + 3i$ and $z_2 = 5 + 9i$ be two complex numbers, then which of the following is

the value of $z_1 + z_2$?

- a) $7 - 12i$
- b) $7 + 12i$
- c) $10 + 27i$
- d) $18 + 15i$

Q

What is roster method to show the set $A = \{x | x \in \mathbb{Z} \wedge 3 < x \leq 3\}$?

- a) $A = \{-3, -2, -1, 0, 1, 2, 3\}$
- b) $A = \{-2, -1, 0, 1, 2, 3\}$
- c) $A = \{-3, -2, -1, 0, 1, 2\}$
- d) $A = \{-3, -2, -1, 1, 2, 3\}$

Q

5.45235452365845125... is a/an

- a) rational number
- b) irrational number
- c) integer
- d) natural number

Q

$$A = \begin{bmatrix} -1 & 2 \\ -3 & -4 \end{bmatrix}$$

Determinant of A is

- a) 12

Created by:

Hafiza Mubeen, Muhammad Nouman, Tasha Khan, M Qamar Nazir Page 3

Mth100 Past papers for mid term

- b) 10
- c) 11
- d) 9

Mth100 Past papers for mid term

Q

Which of the following is Arithmetic Sequence?

- a) $x + a, x + 3a, x + 5a, \dots$
- b) $(x + a) + (x + 3a) + (x + 5a) + \dots$
- c) $(x + a), 2(x + 3a), 4(x + 5a), \dots$
- d) $(x + a) + 2(x + 3a) + 4(x + 5a) + \dots$

Q

$$\begin{bmatrix} 9 & -2 & 4 \\ 5 & 0 & 6 \\ 1 & 3 & 8 \end{bmatrix} - \begin{bmatrix} 4 & 0 & 7 \\ 1 & 5 & -4 \\ -2 & 3 & 2 \end{bmatrix}$$

1) $\begin{bmatrix} 5 & -2 & -3 \\ 4 & -5 & 10 \\ 3 & 0 & 6 \end{bmatrix}$

2) $\begin{bmatrix} 5 & 2 & -3 \\ 4 & -5 & 10 \\ 3 & 0 & 6 \end{bmatrix}$

3) $\begin{bmatrix} 5 & -2 & -3 \\ 4 & 5 & 10 \\ 3 & 0 & 6 \end{bmatrix}$

Q

A line AB has coordinates as A (5, 8) and B (3, 6). The midpoint of AB has coordinates

- a) (1, 1)
- b) (4, 7)
- c) (6.5, 4.5)
- d) (-1.5, -1.5)

Created by:

Hafiza Mubeen, Muhammad Nouman, Tasha Khan, M Qamar Nazir Page 2

Mth100 Past papers for mid term

Q4

Let $z_1 = 3 - 5i$ and $z_2 = -4 - 2i$ be two complex numbers, then which of the following is