



DOWNLOAD SOLVED FINAL

PAST PAPERS BY WAQAR SIDDHU

More in PDF From

VU Answer

Get All Solutions.

Which statement about the General Least Square Method is true?

Answer (Please select your correct option)

VuAnswers.com

Solution obtained by this method is always unique.

This is a numerical method for the solution of System of Linear Equations.

This method find an x that makes Ax as close as possible to the b .

This method gives us exact solution of the system.

*correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell : 03228043306*

Made by: Waqar Siddhu

Let $v = (1, -2, 2, 0)$. The unit vector in the same direction as v is

Answer (Please select your correct option)

VuAnswers.com

$\left(\frac{1}{3}, \frac{2}{3}, \frac{2}{3}, 0\right)$

$\left(\frac{1}{3}, -\frac{2}{3}, \frac{2}{3}, 0\right)$

correct

$\left(\frac{-1}{3}, \frac{2}{3}, \frac{-2}{3}, 0\right)$

$\left(\frac{1}{3}, -\frac{2}{3}, \frac{2}{3}, \frac{1}{3}\right)$

Made by: Waqar Siddhu

Let $u = (3, -2), v = (4, 5)$. For the weighted Euclidean inner product $\langle u, v \rangle = 4u_1v_1 + 5u_2v_2$
 $\langle v, u \rangle =$

Answer (Please select your correct option)

VuAnswers.com

2

-2

3

-3

Correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell no: 03228043306

Made by: Waqar Siddhu

Let $v = (0, 2, 2, 1)$. The unit vector in the same direction as v is

Answer (Please select your correct option)

VuAnswers.com

$\left(0, \frac{2}{3}, \frac{2}{3}, \frac{-1}{3}\right)$

$\left(0, \frac{-2}{3}, \frac{2}{3}, \frac{-1}{3}\right)$

$\left(0, \frac{2}{3}, \frac{2}{3}, \frac{1}{3}\right)$

$\left(0, \frac{-2}{3}, \frac{2}{3}, \frac{1}{3}\right)$

Correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell no: 03228043306

Made by: Waqar Siddhu

Let \mathbb{R}^3 have the Euclidean inner product. Then $u = (2, 1, 3), v = (1, 7, k)$ are orthogonal for

VuAnswers.com

Answer (Please select your correct option)

$k = 9$

$k = -3$

$k = -9$

$k = 3$

Made by: Waqar Siddhu

Let A be $n \times n$ matrix whose entries are real. If λ is an eigenvalue of A with x a corresponding eigenvector in \mathbb{R}^n , then

VuAnswers.com

Answer (Please select your correct option)

$A\bar{x} = \lambda\bar{x}$

$A\bar{x} = \bar{\lambda}\bar{x}$

$Ax = \bar{\lambda}x$

$A\bar{x} = \lambda^{-1}\bar{x}$

Made by: Waqar Siddhu

Suppose that $A = \begin{bmatrix} 1.25 & -0.75 \\ -0.75 & 1.25 \end{bmatrix}$ has eigenvalues 2 and 0.5 .Then origin is a

VuAnswers.com

Answer (Please select your correct option)

Saddle point

correct

correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell : 03228043306

Repellor

Attractor

Made by: Waqar Siddhu

Suppose that $A = \begin{bmatrix} 0.5 & 0.6 \\ -0.3 & 1.4 \end{bmatrix}$ has eigenvalues 0.8 and 1.1 .Then origin is a

VuAnswers.com

Answer (Please select your correct option)

Saddle point

correct

Repellor

Attractor

Made by: Waqar Siddhu

If A is an $m \times n$ matrix with linearly independent column vectors, then A can be factored as

$$A = QR$$

Where Q is an $m \times n$ matrix with orthonormal column vectors, and R is an $n \times n$

VuAnswers.com

Answer (Please select your correct option)

Upper triangular matrix

Invertible matrix

Invertible lower triangular matrix

Invertible upper triangular matrix

Made by: Waqar Siddhu

The matrix equation $A^T A \hat{x} = A^T b$ represents a system of linear equations commonly referred to as the

VuAnswers.com

Answer (Please select your correct option)

normal equations for x

normal equations for \hat{x}

normal equations for A

normal equations for b

Correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell no: 03228043306

Made by: Waqar Siddhu

By the Best Approximation Theorem, the distance from y to W is $\|y - \hat{y}\|$, where $\hat{y} =$

Answer (Please select your correct option)

VuAnswers.com

$\text{proj}_W \hat{y}$

$\text{proj}_W y$

correct

*correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell : 03228043306*

$\text{proj}_y W$

Made by: Waqar Siddhu

$\|u + v + w\| \leq \|u\| + \|v\| + \|w\|$ for all vectors u, v and w in an inner product space.

Answer (Please select your correct option)

VuAnswers.com

True



correct

False



Made by: Waqar Siddhu

The dominant eigenvalue for the matrix $A = \begin{bmatrix} 0 & 0 & 2 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & -3 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$ is

Answer (Please select your correct option)

VuAnswers.com

$\lambda = 1$

$\lambda = -3$

correct

*correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell : 03228043306*

$\lambda = -1$

$\lambda = 0$

Made by: Waqar Siddhu

A square matrix A is invertible if and only if $x = 0$ is not an eigen value of A .

Answer (Please select your correct option)

VuAnswers.com

True



correct

False



Made by: Waqar Siddhu

A square matrix with orthogonal columns _____ matrix. (Click on most appropriate)

VuAnswers.com

Answer (Please select your correct option)

is an orthogonal

Correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell no: 03228043306

may be an orthogonal

may not be an orthogonal

is not an orthogonal

Made by: Waqar Siddhu

If two rows are orthogonal, they are _____.

Answer (Please select your correct option)

VuAnswers.com

linearly independent

Correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell no: 03228043306

linearly dependent

Made by: Waqar Siddhu

If x is orthogonal to both u and v , then x must be _____ to $u + v$.

VuAnswers.com

Answer (Please select your correct option)

orthogonal

orthonormal

perpendicular

parallel

Correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell no: 03228043306

check

Made by: Waqar Siddhu

The given system $2x + 3y = 3$ has
 $6x + 9y = 7$

VuAnswers.com

Answer (Please select your correct option)

Unique solution

Infinitely many solutions

No solution

None of these

Correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell no: 03228043306

Made by: Waqar Siddhu

Which statement about the matrix $\begin{bmatrix} 1 & 0 & 0 & 0 \\ 7 & 2 & 0 & 0 \\ 9 & 1 & 2 & 0 \\ 5 & 4 & 2 & -1 \end{bmatrix}$ is false?

Answer (Please select your correct option)

VuAnswers.com

Eigenvalue 2 has Algebraic multiplicity 1

Eigenvalue of the matrix are 1, 2 and -1.

Characteristic polynomial of the matrix is $(1-\lambda)(2-\lambda)^2(-1-\lambda)$.

Eigenvalue -1 has multiplicity 1.



Made by: Waqar Siddhu

If $A = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}$ is diagonalizable then A has 2 distinct eigenvalues.

Answer (Please select your correct option)

VuAnswers.com

True



0 1
1 0

Correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell no: 03228043306

False



Made by: Waqar Siddhu

A is diagonalizable if $A = PDP^{-1}$ Where

Answer (Please select your correct option)

VuAnswers.com

D is any matrix and P is an invertible matrix

D is a diagonal matrix and P is any matrix

D is a diagonal matrix and P is invertible matrix

D is a invertible matrix and P is any matrix

Correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell no: 03228043306

Made by: Waqar Siddhu

Which statement is FALSE.

Answer (Please select your correct option)

VuAnswers.com

If $Ax = \lambda x$ for some real number λ then λ is known as eigenvalue of the matrix A .

The eigenvalues of any matrix are on its main diagonal.

In order to find the eigenvalues we solve the equation $|A - \lambda I| = 0$

An eigenspaces of A is the Null space of some matrix.

Made by: Waqar Siddhu

If a set $S = \{1, x, x^2\}$ is a basis for p_2 and $[p]_S = (2, 4, 7)$, then which of the following is the most appropriate option ?

Answer (Please select your correct option)

VuAnswers.com

$p_2 = 2 - 4x + 7x^2$

$p_2 = 2 - 4x - 7x^2$

$p_2 = 2 + 4x + 7x^2$

$p_2 = 4x - 7x^2$

Correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell no: 03228043306

Made by: Waqar Siddhu

Which of the following is the set of standard basis for R^3 ?

Answer (Please select your correct option)

VuAnswers.com

$\{(1, 1, 0), (0, 1, 0), (1, 0, 1)\}$

$\{(1, 0, 0), (0, 1, 0), (0, 0, 1)\}$

$\{(1, 0, 0), (1, 1, 0), (0, 0, 1)\}$

$\{(1, 0, 0), (0, 1, 0), (1, 1, 1)\}$

Correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell no: 03228043306

Made by: Waqar Siddhu

Consider the bases for R^2 given by $B = \{ \overset{\uparrow}{b}_1, \overset{\uparrow}{b}_2 \}$ and $C = \{ \overset{\uparrow}{c}_1, \overset{\uparrow}{c}_2 \}$; where $\overset{\uparrow}{b}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$, $\overset{\uparrow}{b}_2 = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$, $\overset{\uparrow}{c}_1 = \begin{bmatrix} -5 \\ 3 \end{bmatrix}$, $\overset{\uparrow}{c}_2 = \begin{bmatrix} -2 \\ 1 \end{bmatrix}$, also assume that $P_{B \leftarrow C} = \begin{bmatrix} -2 & -1 \\ 3 & 1 \end{bmatrix}$; then which of the following is the change-of-coordinates matrix from B to C ?

Answer (Please select your correct option)

VuAnswers.com

$P_{C \leftarrow B} = \begin{bmatrix} 1 & 1 \\ -3 & -2 \end{bmatrix}$

$P_{C \leftarrow B} = \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}$

$P_{C \leftarrow B} = \begin{bmatrix} -5 & -2 \\ 3 & 1 \end{bmatrix}$

$P_{C \leftarrow B} = \begin{bmatrix} -8 & -3 \\ 3 & 1 \end{bmatrix}$

Correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell no: 03228043306

Made by: Waqar Siddhu

If the general term of a typical signal is $(0.6)^k$, then determine which of the following is the signal for $k = -2$?

Answer (Please select your correct option)

VuAnswers.com

$(0.6)^{-2} = 0$

$(0.6)^{-2} = 0.6$

$(0.6)^{-2} = (0.6)^2$

$(0.6)^{-2} = 1/(0.6)^2$

Correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell no: 03228043305

Made by: Waqar Siddhu

If the Casorati matrix is not invertible , then which of the following is the most appropriate option regarding the associated signals ?

Answer (Please select your correct option)

VuAnswers.com

The signals are linearly independent .

Correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell no: 03228043306

The signals are linearly dependent .

The signals may or may not dependent .

The signals may or may not independent .

Made by: Waqar Siddhu

If $\{y_k\} = \{\dots, 1, 0.7, 0, -0.7, -1, -0.7, 0, 0.7, 1, 0.7, 0, \dots\}$ and $0.35y_{k+2} + 0.6y_{k+1} + 0.42y_k = z_k$;

↑

$k = 0$

then which of the following is the value of z_0 ?

VuAnswers.com

Answer (Please select your correct option)

0.840

0.049

-0.770

- 1.139

Correct answer solved by Hadi

Email : usmanraj20@gmail.com

Cell no: 03228043306

Made by: Waqar Siddhu

Suppose that $B = \{b_1, b_2\}$ is a basis for V and $C = \{c_1, c_2, c_3\}$ is a basis for W . Let $T: V \rightarrow W$ be a linear transformation with the property that $T(b_1) = 5c_1 - 2c_2 + 3c_3$ and $T(b_2) = 4c_1 - c_2 + 7c_3$. Determine the value of $[T(b_2)]_C$?

Answer (Please select your correct option)

VuAnswers.com

$\begin{bmatrix} 5 \\ -2 \\ 3 \end{bmatrix}$

$\begin{bmatrix} 7 \\ -1 \\ 7 \end{bmatrix}$

$\begin{bmatrix} 4 \\ -1 \\ 7 \end{bmatrix}$

$\begin{bmatrix} 3 \\ -2 \\ 7 \end{bmatrix}$

Made by: Waqar Siddhu

Determine whether the set of vectors are orthogonal or not.

$$\begin{bmatrix} 5 \\ -4 \\ 0 \end{bmatrix}, \begin{bmatrix} -4 \\ 1 \\ -3 \end{bmatrix}, \begin{bmatrix} 3 \\ 3 \\ 5 \end{bmatrix}$$

Answer ([Please click here to Add Answer](#))

VuAnswers.com



Normal Arial 12 B I U

Made by: Waqar Siddhu

Question No : 1 of 52

Marks: 1 (Budgeted Time 1 Min)

Which statement about the General Least Square Method is true?

Answer (Please select your correct option)

- Solution obtained by this method is always unique.
- This is a numerical method for the solution of System of Linear Equations.
- This method find an x that makes Ax as close as possible to the b .
- This method gives us exact solution of the system.

Correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell no: 03228043306

Start Time: 11:33 PM

120:00
Time Left

1



Question No : 2 of 52

Marks: 1 (Budgeted Time 1 Min)

Let $v = (1, -2, 2, 0)$. The unit vector in the same direction as v is

Answer (Please select your correct option)

$\left(\frac{1}{3}, \frac{2}{3}, \frac{2}{3}, 0\right)$

$\left(\frac{1}{3}, \frac{-2}{3}, \frac{2}{3}, 0\right)$

ans

$\left(\frac{-1}{3}, \frac{2}{3}, \frac{-2}{3}, 0\right)$

$\left(\frac{1}{3}, \frac{-2}{3}, \frac{2}{3}, \frac{1}{3}\right)$

Correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell no: 03228043306

Start Time: 11:33 PM

120:00

Time Left



2



Question No : 3 of 52

Marks: 1 (Budgeted Time 1 Min)

Let $v = (1, -2, 2, 0)$. The unit vector in the same direction as v has magnitude

Answer (Please select your correct option)

 3

ans

Correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell no: 03228043306

 1 2 -1

Start Time: 11:33 PM

120:00

Time Left



3



Question No : 4 of 52

Marks: 1 (Budgeted Time 1 Min)

Let \mathbb{R}^3 have the Euclidean inner product. Then $u = (2, 1, 3), v = (1, 7, k)$ are orthogonal for

out of

Answer (Please select your correct option)

 $k = 9$ $k = -3$ $k = -9$ $k = 3$

Start Time: 11:33 PM

120:00
Time Left

4



Question No : 5 of 52

Marks: 1 (Budgeted Time 1 Min)

Let \mathbb{R}^3 have the Euclidean inner product. Then $u = (k, -3, 1), v = (-3, 5, 6)$ are orthogonal for

out of

Answer (Please select your correct option)

$k = 2$

$k = -2$

$k = 3$

$k = -3$

Start Time: 11:33 PM

120:00
Time Left

5



Question No : 6 of 52

Marks: 1 (Budgeted Time 1 Min)

If u and v are orthogonal then

Answer (Please select your correct option)

$\|u - v\|^2 = \|u\| + \|v\|$

$\|u - v\| = \|u\|^2 + \|v\|^2$

$\|u - v\|^2 = \|u\|^2 + \|v\|^2$

$\|u - v\|^{\frac{1}{2}} = \|u\| + \|v\|$


Start Time: 11:33 PM

120:00

Time Left



6



Question No : 7 of 52

Marks: 1 (Budgeted Time 1 Min)

Suppose that $A = \begin{bmatrix} 0.5 & 0.6 \\ -0.3 & 1.4 \end{bmatrix}$ has eigenvalues 0.8 and 1.1. Then origin is a

Answer (Please select your correct option)

 Saddle point

Correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell no: 03228043306

 Repellor Attractor

Start Time: 11:33 PM

120:00

Time Left



7



Question No : 7 of 52

Marks: 1 (Budgeted Time 1 Min)

Suppose that $A = \begin{bmatrix} 0.5 & 0.6 \\ -0.3 & 1.4 \end{bmatrix}$ has eigenvalues 0.8 and 1.1. Then origin is a

Answer (Please select your correct option)

 Saddle point

Correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell no: 03228043306

 Repellor Attractor

Start Time: 11:33 PM

119:00

Time Left



7



Question No : 8 of 52

Marks: 1 (Budgeted Time 1 Min)

If A is an $m \times n$ matrix with linearly independent column vectors, then A can be factored as

$$A = QR$$

Where Q is an $m \times n$ matrix with orthonormal column vectors, and R is an $n \times n$

Answer (Please select your correct option)

 Upper triangular matrix

ans

 Invertible matrix Invertible lower triangular matrix Invertible upper triangular matrix

Start Time: 11:33 PM

119:00
Time Left

8



Question No : 9 of 52 **Marks: 1 (Budgeted Time 1 Min)**

The QR-Decomposition of a 3×3 matrix A gives

Answer (Please select your correct option)

- matrix Q of order 3×1 and R of order 3×1
- matrix Q of order 3×3 and R of order 3×1
- matrix Q of order 3×3 and R of order 3×3

ans

1 0 0
0 1 0
0 0 1

Correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell no: 03228043306

Start Time: 11:33 PM
119:00
Time Left



9

Information, Previous, Next, Previous, Next, Print icons

Question No : 10 of 52

Marks: 1 (Budgeted Time 1 Min)

The QR-Decomposition of a 5×2 matrix A gives

Answer (Please select your correct option)

 matrix Q of order 5×1 and R of order 5×1 matrix Q of order 5×2 and R of order 2×5 matrix Q of order 5×2 and R of order 2×2

Start Time: 11:33 PM

119:00

Time Left



10



Question No : 11 of 52

Marks: 1 (Budgeted Time 1 Min)

If u, v and w are vectors in a real inner product space, and k is any scalar, then which one of the following is not a property for the inner product space

out of

Answer (Please select your correct option)

$\langle 0, v \rangle = \langle v, 0 \rangle$

$\langle u, kv \rangle = k \langle u, v \rangle$

$\langle u, v - w \rangle = \langle u, v \rangle + \langle u, w \rangle$

Start Time: 11:33 PM

119:00

Time Left



11



Question No : 12 of 52

Marks: 1 (Budgeted Time 1 Min)

The dominant eigenvalue for the matrix $A = \begin{bmatrix} 0 & 0 & 2 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & -3 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$ is

Answer (Please select your correct option)

 $\lambda = 1$ $\lambda = -3$

ans

 $\lambda = -1$ $\lambda = 0$

Correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell no: 03228043306

Start Time: 11:33 PM

119:00

Time Left



12



Question No : 13 of 52 **Marks: 1 (Budgeted Time 1 Min)**

_____ of a matrix is the sum of main diagonal elements of that matrix.

Answer (Please select your correct option)

- Trace
- Determinent
- Eigenvalue
- Sum

ans

Start Time: 11:33 PM
119:00
Time Left

13

Navigation icons: Information, Previous, Home, Next, Last

Question No : 14 of 52 **Marks: 1 (Budgeted Time 1 Min)**

If a square matrix has orthonormal columns, then it also has _____.

Answer (Please select your correct option)

- orthonormal rows
- ans**
- orthonormal diagonal

Correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell no: 03228043306

Start Time: 11:33 PM
119:00

14



Question No : 15 of 52

Marks: 1 (Budgeted Time 1 Min)

If x is orthogonal to both u and v , then x must be _____ to $u - v$.

Answer (Please select your correct option)

 orthogonal orthonormal perpendicular parallel


Start Time: 11:33 PM

119:00

Time Left



15



Question No : 16 of 52

Marks: 1 (Budgeted Time 1 Min)

If a matrix U has orthonormal columns, then _____ = I

Answer (Please select your correct option)

 UU^T

ans

 UU U^{-1} $U+U^T$

Start Time: 11:33 PM

119:00

Time Left



16



Question No : 17 of 52

Marks: 1 (Budgeted Time 1 Min)

If 5 is an eigenvalue of A and x is a corresponding eigenvector, then the eigenvalue of A^2 is

Answer (Please select your correct option)

- 5
- 25
- 10
- 15

ans

$$A^2 = 5(2) = 25$$

Correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell no: 03228043306


Start Time: 11:33 PM

119:00

Time Left



17



Question No : 18 of 52 Marks: 1 (Budgeted Time 1 Min)

The eigenvalues of the matrix $\begin{bmatrix} 3 & -4 & 1 \\ 0 & -5 & 1 \\ 0 & 0 & 4 \end{bmatrix}$ are

we read down to up

Answer (Please select your correct option)

- 3,-4,1
- 4,-5,3
- 3,-5,1
- 4,1,-5

ans

Correct answer solved by Hadi
 Email : usmanraj20@gmail.com
 Cell no: 03228043306

$$\begin{bmatrix} 3 & & \\ & -5 & \\ & & 4 \end{bmatrix}$$

Start Time: 11:33 PM
119:00
 Time Left



18




Question No : 19 of 52

Marks: 1 (Budgeted Time 1 Min)

If $A = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}$ is diagonalizable, then A is invertible.

Answer (Please select your correct option)

True

 ans

False

Start Time: 11:33 PM

119:00

19



Question No : 20 of 52

Marks: 1 (Budgeted Time 1 Min)

An $n \times n$ matrix A is diagonalizable if and only if A has

Answer (Please select your correct option)

 n linearly independent eigenvectors ans n^2 linearly independent eigenvectors $n + 1$ linearly independent eigenvectors

Correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell no: 03228043306

Start Time: 11:33 PM

119:00
Time Left

20



Question No : 21 of 52

Marks: 1 (Budgeted Time 1 Min)

Let A have eigenvalues 2, 5, 0, -7, and -2. Then the dominant eigenvalue for A is

Answer (Please select your correct option)

 $\lambda = 5$ $\lambda = 0$ $\lambda = -7$ $\lambda = 2$

Correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell no: 03228043306

Start Time: 11:33 PM

119:00
Time Left

21



Question No : 22 of 52

Marks: 1 (Budgeted Time 1 Min)

Which statement is FALSE.

Answer (Please select your correct option)

If $Ax = \lambda x$ for some real number λ then λ is known as eigenvalue of the matrix A.

The eigenvalues of any matrix are on its main diagonal.

ans

In order to find the eigenvalues we solve the equation $|A - \lambda I| = 0$

An eigenspaces of A is the Null space of some matrix.

Start Time: 11:33 PM

119:00

Time Left



22



Question No : 23 of 52 **Marks: 1 (Budgeted Time 1 Min)**

Algebra is a transformation of real life problems into sort of _____.

Answer (Please select your correct option)

- logical representation
- mathematical representation
- physical representation
- illogical representation

ans

Correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell no: 03228043306

Start Time: 11:33 PM
119:00
Time Left

23

Navigation icons: Information, Previous, Home, Next, Last

Question No : 24 of 52

Marks: 1 (Budgeted Time 1 Min)

If reduced echelon form of a linear system is $\begin{bmatrix} 1 & 0 & 5 & 5 \\ 0 & 1 & 1 & 6 \\ 0 & 0 & 0 & 0 \end{bmatrix}$ when free variable $x_3 = 0$, then which of the following is true for it?

Answer (Please select your correct option)

 The particular solution is $(0, 5, 6)$. The particular solution is $(6, 5, 0)$. The particular solution is $(5, 6, 0)$. The particular solution is $(0, 6, 5)$.

ans

Start Time: 11:33 PM

119:00

Time Left



24



Question No : 25 of 52

Marks: 1 (Budgeted Time 1 Min)

If the vector equation $c_1 \vec{v}_1 + c_2 \vec{v}_2 = 0$ with $c_1 = 0 = c_2$ then which of the following is true for $\{\vec{v}_1, \vec{v}_2\}$?

Answer (Please select your correct option)

 It is a linearly independent set.

ans

Correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell no: 03228043306

 It is a linearly dependent set. The system of equations is inconsistent. The system of equations is non- homogeneous.


Start Time: 11:33 PM

119:00

Time Left



25



Question No : 26 of 52

Marks: 1 (Budgeted Time 1 Min)

Which of the following is true for the linear operator L defined by $L\left(\begin{bmatrix} a_1 \\ a_2 \end{bmatrix}\right) = \begin{bmatrix} a_1 \\ -a_2 \end{bmatrix}$?

Answer (Please select your correct option)

 It is an enlargement by a negative scale factor. It is a shear.

ans

 It is a reflection about X -axis. It is a reflection about Y -axis.

Correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell no: 03228043306

Start Time: 11:33 PM

119:00

Time Left



26



Question No : 27 of 52

Marks: 1 (Budgeted Time 1 Min)

If $A = \begin{bmatrix} 4 & -1 \\ 5 & 3 \end{bmatrix}$, then which of the following is the value of $\det(A)$?

Answer (Please select your correct option)

 7 -17 17 11

ans

$$4(3) - (-1)(5) = 17$$

Correct answer solved by Hadi

Email : usmanraj20@gmail.com

Cell no: 03228043306


Start Time: 11:33 PM

119:00

Time Left



27



Question No : 28 of 52

Marks: 1 (Budgeted Time 1 Min)

If the matrix $A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 2 \\ 0 & 0 & 2 \end{bmatrix}$, then which of the following is true about it ?

Answer (Please select your correct option)

 Its determinant is 0 . Its determinant is 1 . Its determinant is 2 .

ans

 Its determinant is 4 .

Start Time: 11:33 PM

119:00

Time Left



28



Question No : 29 of 52

Marks: 1 (Budgeted Time 1 Min)

If the matrix $A = \begin{bmatrix} 1 & 5 & 4 \\ 0 & 1 & 7 \\ 0 & 0 & 0 \end{bmatrix}$, then which of the following is true about it ?

Answer (Please select your correct option)

 Its determinant is 0 .

ans

Correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell no: 03228043306

 Its determinant is 2 . Its determinant is 4 . Its determinant is 6 .

Start Time: 11:33 PM

119:00

Time Left



29



Question No : 30 of 52

Marks: 1 (Budgeted Time 1 Min)

If the determinant of the matrix $A = \begin{bmatrix} 4 & 3 & 5 \\ 3 & 1 & 1 \\ 5 & 7 & 7 \end{bmatrix}$ is 32 and the matrix B is obtained by multiplying any row of A with an integer value 4, then which of the following is true about the matrix B ?

Answer (Please select your correct option)

 Its determinant is 18 . Its determinant is - 32 . Its determinant is 128 .

i think

 The information is not sufficient to calculate the determinant .

Start Time: 11:33 PM

119:00

Time Left



30



Question No : 31 of 52

Marks: 1 (Budgeted Time 1 Min)

If a set W be a subspace of a vector space V , then which of the following is NOT true for it ?

Answer (Please select your correct option)

 It must be closed under the scalar multiplication . It may or may not be closed under the operation of addition . It must have an additive inverse of each element . It must be commutative under the operation of addition .

ans

Correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell no: 03228043306

Start Time: 11:33 PM
119:00
Time Left

31



Question No : 32 of 52

Marks: 1 (Budgeted Time 1 Min)

Let a set S is a basis of a vector space V , then which of the following is NOT true about it ?

Answer (Please select your correct option)

 It spans V . It is linearly independent. It is linearly dependent.

ans

 Each element of S belongs to V .

Correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell no: 03228043306


Start Time: 11:33 PM

119:00

Time Left



32



Question No : 33 of 52

Marks: 1 (Budgeted Time 1 Min)

If a set $S = \left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$ is a basis for R^3 and $[\mathbf{v}]_S^r = \begin{bmatrix} 7 \\ 5 \\ 3 \end{bmatrix}$, then which of the following is the most appropriate option ?

Answer (Please select your correct option)

$\mathbf{v}^r = (1, 0, 0)^t \in R^3$

$\mathbf{v}^r = (2, 3, 4)^t \in R^3$

$\mathbf{v}^r = (7, 5, 3)^t \in R^3$

$\mathbf{v}^r = (2, 7, 4)^t \in R^3$

ans

Correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell no: 03228043306

Start Time: 11:33 PM

119:00

Time Left



33



Question No : 34 of 52

Marks: 1 (Budgeted Time 1 Min)

If P is a polynomial of order ' n ', then which of the following is true for the $\dim(P_n)$?

Answer (Please select your correct option)

 $\dim(P_n) = n - 1$ $\dim(P_n) = n$ $\dim(P_n) = n + 1$ $\dim(P_n) = 1$

Correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell no: 03228043306

Start Time: 11:33 PM

119:00
Time Left

34



Question No : 35 of 52

Marks: 1 (Budgeted Time 1 Min)

Let $A = \begin{bmatrix} 1 & 3 & 2 & -1 & 5 \\ 2 & 2 & 1 & 1 & 4 \\ -1 & 4 & 1 & 2 & -3 \\ 3 & 5 & 3 & -4 & 2 \end{bmatrix}$, the row space of A is the subspace of R^5 . Identify the set by which the row space of A is spanned?

Answer (Please select your correct option)

 $\{(1, 2, -1, 3), (2, 4, -2, 6), (2, 1, 1, 3), (-1, 1, 2, -4), (6, 3, 3, 9)\}$ $\{(1, 3, 2, -1, 5), (2, 6, 4, -2, 10), (-1, 4, 1, 2, -3), (-3, 12, 3, 6, -9)\}$ $\{(1, 3, 2, -1, 5), (2, 2, 1, 1, 4), (-1, 4, 1, 2, -3), (3, 5, 3, -4, 2)\}$ $\{(1, 2, -1, 3), (3, 2, 4, 5), (2, 1, 1, 3), (-1, 1, 2, -4), (5, 4, -3, 2)\}$

ans

Start Time: 11:33 PM

119:00

Time Left



35



Question No : 36 of 52

Marks: 1 (Budgeted Time 1 Min)

If $A = \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix}$ and $\vec{x} \in \text{Row}(A)$, then which of the following is the most appropriate option ?

Answer (Please select your correct option)

$\vec{x} = c_1(1, 3) + c_2(2, 4)$

$\vec{x} = c_1(1, 2) + c_2(3, 4)$

ans

Correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell no: 03228043306

$\vec{x} = c_1(1, 4) + c_2(3, 2)$

$\vec{x} = c_1(2, 3) + c_2(4, 1)$

Start Time: 11:33 PM

119:00

Time Left



36



Question No : 37 of 52

Marks: 1 (Budgeted Time 1 Min)

Consider the bases for R^2 given by $B = \{ \overset{\uparrow}{b}_1, \overset{\uparrow}{b}_2 \}$ and $C = \{ \overset{\uparrow}{c}_1, \overset{\uparrow}{c}_2 \}$; where $\overset{\uparrow}{b}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$, $\overset{\uparrow}{b}_2 = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$, $\overset{\uparrow}{c}_1 = \begin{bmatrix} -6 \\ 4 \end{bmatrix}$, $\overset{\uparrow}{c}_2 = \begin{bmatrix} -4 \\ 7 \end{bmatrix}$; then which of the following is the change-of-coordinates matrix from C to B ?

Answer (Please select your correct option)

$P_{B \leftarrow C} = \begin{bmatrix} -2 & 3 \\ 4 & 7 \end{bmatrix}$

$P_{B \leftarrow C} = \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}$

$P_{B \leftarrow C} = \begin{bmatrix} -6 & -4 \\ 4 & 7 \end{bmatrix}$

$P_{B \leftarrow C} = \begin{bmatrix} -10 & -11 \\ 4 & 7 \end{bmatrix}$

ans

Correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell no: 03228043306

Start Time: 11:33 PM

119:00

Time Left



37



Question No : 38 of 52

Marks: 1 (Budgeted Time 1 Min)

If $B = \{ \hat{b}_1, \hat{b}_2, \dots, \hat{b}_n \}$ and E is the standard basis $\{ e_1, e_2, \dots, e_n \}$ in R^n ; then which of the following is the value of $\left[\hat{b}_1 \right]_E$?

Answer (Please select your correct option)

 E $\hat{b}_1 + E$ \hat{b}_1 \hat{b}_2

Start Time: 11:33 PM

119:00
Time Left

38



Question No : 39 of 52

Marks: 1 (Budgeted Time 1 Min)

If $\{y_k\} = \{\dots, 1, 0.7, 0, -0.7, -1, -0.7, 0, 0.7, 1, 0.7, 0, \dots\}$ and $0.35y_{k+2} + 0.6y_{k+1} + 0.42y_k = z_k$;

 \uparrow $k = 0$

then which of the following is the value of z_0 ?

Answer (Please select your correct option)

 0.840 0.049 -0.770 -1.139

Correct answer solved by Hadi
Email : usmanraj20@gmail.com
Cell no: 03228043306

ans


Start Time: 11:33 PM

119:00

Time Left



39



MORE PAST PAPERS BY WAQAR SIDDHU

Provide Solved in PDF From

VU Answer

Get All Solutions.

